

改訂コルポスコピー所見分類: 日本婦人科腫瘍学会2014

A) 総合評価 General assessment	GA
1. 観察可 観察不可 (理由: 炎症、出血、瘢痕など) Adequate or inadequate for the reason (inflammation, bleeding, scar, etc)	ADE or INA
2. 扁平円柱境界 Squamocolumnar junction	SCJ
可視 Completely visible	V1
部分的可視 Partially visible	V2
不可視 Not visible	V3
3. 移行帯 Transformation zone	TZ
1型 Type 1	TZ1
2型 Type 2	TZ2
3型 Type 3	TZ3
B) 正常所見 Normal colposcopic findings	NCF
1. 扁平上皮 Original squamous epithelium	S
2. 円柱上皮 Columnar epithelium	C
3. 化生上皮 Metaplastic squamous epithelium	T
ナボット卵 Nabothian cysts	N
腺開口 Gland openings	Go
C) 異常所見 Abnormal colposcopic findings	ACF
1. 概観 General principles	
病変の部位: 移行帯 (内、外) (時方向) Location of the lesion: inside or outside the transformation zone (clock position)	
病変の大きさ: 子宮腔部占拠率 (%) Size of the lesion: percentage of cervix the lesion covers	
2. 軽度所見 Grade 1 (minor)	
白色上皮 (軽度) Thin acetowhite epithelium	W1
モザイク (軽度) Fine mosaic	M1
赤点斑 (軽度) Fine punctation	P1
不規則・地図状辺縁 Irregular, Geographic border	B1
3. 高度所見 Grade 2 (major)	
白色上皮 (高度) Dense acetowhite epithelium	W2
モザイク (高度) Coarse mosaic	M2
赤点斑 (高度) Coarse punctation	P2
異常腺開口 Abnormal gland openings	aGo
鋭角辺縁、内部境界、尾根状隆起 Sharp border, Inner border, Ridge sign	B2
4. 非特異的所見 Nonspecific findings	
白斑 (角化、過角化) Leukoplakia (keratosis, hyperkeratosis)	L
びらん Erosion	Er
D) 浸潤癌所見 Suspicious for invasion	IC
異型血管 Atypical vessels	aV
付随所見 Additional signs: fragile vessels, irregular surface, exophytic lesion, necrosis, ulceration (necrotic), tumor or gross neoplasm	
E) その他の非癌所見 Miscellaneous findings	MF
1. コンジローマ Condyloma	Con
2. 炎症 Inflammation	Inf
3. 萎縮 Atrophy	Atr
4. ポリープ (頸管外、頸管内) Polyp (ectocervical or endocervical)	Po
5. 潰瘍 Ulcer	Ul
6. その他 Others	etc